#### **ABSCESS PREVENTION**

#### Avoiding abscesses

- Wash your hands and the injection site.
- Use alcohol pads and wipe back & forth (rub hard) over injection site to remove dirt.
- Then use another new alcohol pad for the final cleaning.

#### What is a skin abscess?

- Pocket of pus
- Often found at injection sites, but can be found elsewhere
- More likely with skin-popping muscling missing a vein
- May occur even after you stop injecting

### How do you know it's an abscess?

- Pink or reddish lump on skin
- Tender or painful
- Warm or hot to touch
- Red streaks and/or rapid swelling indicates infection is spreading

# How can an abscess make you feel?

- Tired or sick
- Feverish, may have chills (continued above)

Chest pains may occur if infection goes to heart or lungs

### What should I do if I get an abscess?

- Treat at home with warm soaks only if:
  - No red streaks
  - Skin not hot and puffy
- Go to a clinic if abscess:
  - Notimproving, especially after 5-7 days
  - Gets bigger and/or very painful
  - Is hot and puffy

Red streaks start spreading from the abscess-go ASAP!

Go to emergency room if:

Chest pain
High fever, chills
Infection looks like it is
spreading fast

### Warning

- Do not shoot into or near an abscess
- Do not squeeze or cut into an abscess yourself - infection could move into blood, then heart/lungs
- Do not leave draining abscesses uncovered the drainage is contagious

## How do you soak/use compresses?

- Use warm/hot water (that doesn't burn your skin)
- Soak in tub of plain hot water or hot water with Epsom salts
- Use hot, wet, clean washcloth and hold on abscess, if abscess cannot be soaked in tub
- Soak abscess 3 to 4 times a day for 10-15 minutes each time, if possible
- Cover with a clean dry bandage after soaking
- STOP soaking/using compresses when abscess starts draining

# What about antibiotics?



- Take all antibiotics, if prescribed, even if you feel better
- Take antibiotics after you fix (if
- using heroin)
- Do not take antibiotics with alcohol, juice, or soda because they may upset your stomach
- Do not share antibiotics because someone else's pills may not cure your infection
- Return to the doctor if your infection is not better after taking antibiotics