

# Toledo-Lucas County Health Department Seasonal Influenza Activity Summary

MMWR Week 06 February 2-8, 2020

## Local Surveillance:

### Current Influenza Activity Levels:

Lucas County had **37** confirmed Influenza-associated Hospitalizations (IAH) reported in week 06, bringing the total to **282** cases as of February 8, 2020. 54% of cases are flu type B, 46% are flu type A (37% reported H1N1, 63% unsubtype)

--There were no deaths reported



Gender Distribution of Cases: 56% female, 44% male

Lucas County accounts for 4.0% of the IAH reported in Ohio. Rate per 100,000 population\*\*: 49.12

\*Stating Influenza as the cause of death OR as 'a consequence of' on the certificate of death. Please note that some individuals may have underlying health conditions and influenza may not be the sole cause of their death.

\*\*Disease rates were calculated by number of cases per 100,000 residents using 2010 census data

Graph 1 illustrates the current confirmed IAH compared to baseline, which is a five year average.




\* Combination of ED visit data and Influenza-associated Hospitalizations.


ED Visits are comprised of Influenza-like illness and respiratory syndrome as compared to the baseline average

# Toledo-Lucas County Health Department Seasonal Influenza Activity Summary

MMWR Week 06 February 2-8, 2020



The number of emergency department visits with patients exhibiting fever/ILI symptoms was slightly above baseline for MMWR 06. ILI (Influenza-Like Illness) is defined as a *fever greater than or equal to 100 degrees F AND a cough or sore throat*. (Graph 3)



**Summary of State Data:** Current Ohio Activity Level (Geographic Spread) – **Widespread** Definition: Increased ILI in at least half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the state. During MMWR Week 6, public health surveillance data sources indicate high intensity for influenza-like illness (ILI) in outpatient settings reported by Ohio's sentinel providers. The percentage of emergency department visits with patients exhibiting constitutional symptoms increased and are above baseline levels statewide; fever and ILI specified ED visits increased and are also above baseline levels. Reported cases of influenza-associated hospitalizations are above the seasonal threshold\*. There were 994 influenza-associated hospitalizations reported during MMWR Week 6.

# Toledo-Lucas County Health Department Seasonal Influenza Activity Summary

MMWR Week 06 February 2-8, 2020

## **State Surveillance:**

- ODH lab has reported 539 positive influenza tests from specimens sent from sentinel ILINet providers and hospital clinical labs. 2019-2020 influenza season results: (263) A/pdmH1N1; (17) A/H3N2; (259) Influenza B; (through 02/08/2020).
- The National Respiratory and Enteric Virus Surveillance System (NREVSS) has reported 54,929 influenza specimens tested by RT-PCR at participating facilities. 2019-2020 influenza season positive results: (262) A/pdmH1N1; (3) A/H3N2; (4,479) Flu A Not Subtyped; and (6,520) Flu B; (through 02/08/2020)
- 2 influenza-associated pediatric mortalities have been reported during the 2019-2020 season (through 02/08/2020).
- No novel influenza A virus infections have been reported during the 2019-2020 season (through 02/08/2020).
- Incidence of confirmed influenza-associated hospitalizations in 2019-2020 season = 5,457 (through 02/08/2020).

**Regional Surveillance\*:** During week 5 (January 26th – February 1st, 2020), the proportion of outpatient visits for ILI in Region 5 (Ohio is in Region 5) was 4.84%, which is above the regional baseline of 1.9%. Michigan, Ohio, West Virginia, Pennsylvania, Kentucky, and Indiana reported Widespread Activity.

**National Surveillance\*:** During week 5 (January 26th – February 1st, 2020), the majority of the U.S. reported High influenza activity. The proportion of outpatient visits for ILI was 6.7%, which is above the national baseline of 2.4%. All 10 HHS regions reported ILI levels at or above their region-specific baseline level. Nationally, influenza B/Victoria viruses have been reported more frequently than other influenza viruses this season. However, during recent weeks, influenza A(H1N1)pdm09 viruses have been reported more frequently than B/Victoria viruses. The predominant virus varies by region.

*\*Regional and National surveillance data are reported one week later than Ohio state and local data*

**State, Regional and National data is provided by the Ohio Department of Health**

[ODH Influenza Activity Summary MMWR 6](#)